

The lowest resonance of QCD

H. Leutwyler

u^b



Novosibirsk, 28 Feb. 2006

$\pi\pi$ interaction

- Plays a crucial role whenever the strong interaction is involved at low energies
Example: Standard model prediction for muon magnetic moment
- Main experiments on $\pi\pi$ scattering were done in the seventies. What's new ?
 - Significant theoretical progress, based on ChPT + dispersion theory
 - New precision data:

$K \rightarrow \pi\pi\ell\nu$	E865	Brookhaven
pionic atoms	DIRAC	CERN
$K \rightarrow 3\pi$	NA48/2	CERN
 - Lattice results on M_π , F_π , a_0^2 , $\langle r^2 \rangle_s$

Analyticity and crossing

- $\pi\pi$ scattering is special: crossed channels are identical
- ⇒ $\text{Re } T(s, t)$ can be represented as a twice subtracted dispersion integral over $\text{Im } T(s, t)$ in physical region

S.M. Roy 1971

- The 2 subtraction constants can be identified with the *S*-wave scattering lengths:

$$a_0^0, \quad a_0^2 \begin{array}{l} \leftarrow \text{isospin} \\ \leftarrow \text{angular momentum} \end{array}$$

- Representation leads to dispersion relations for the individual partial waves: *Roy equations*

Roy equations

- Pioneering work on the physics of the Roy equations:
Basdevant, Froggatt & Petersen 1974
- Dispersion integrals converge rapidly (2 subtractions)
 - ⇒ Crude phenomenological information on $\text{Im } T(s, t)$ for energies above 800 MeV suffices
 - ⇒ Given a_0^0, a_0^2 , the scattering amplitude can be calculated to within small uncertainties


Ananthanarayan, Colangelo, Gasser & L. 2001
Descotes, Fuchs, Girlanda & Stern 2002

- ⇒ a_0^0, a_0^2 are the essential parameters at low energy
- Main problem in early work: a_0^0, a_0^2 poorly known
Experimental information near threshold is meagre

Low energy theorems


- Chiral perturbation theory provides the missing piece: theoretical prediction for a_0^0, a_0^2
Weinberg 1966, Gasser & L. 1984, Bijnens, Colangelo, Ecker, Gasser & Sainio 1996
- Most accurate results for a_0^0, a_0^2 are obtained by matching the chiral and dispersive representations near the center of the Mandelstam triangle
Colangelo, Gasser & L. 2001
- In combination with the low energy theorems for a_0^0, a_0^2 , the dispersion relations for the partial waves fix the $\pi\pi$ scattering amplitude to an incredible degree of accuracy

Predictions for the S-wave $\pi\pi$ scattering lengths



Sizeable corrections in a_0^0 , while a_0^2 nearly stays put

Tests of the predictions for a_0^0 , a_0^2 : experiment & lattice



Theory is ahead of experiment . . .

The σ

Work done in collaboration with I. Caprini and G. Colangelo, hep-ph/0512364

- Does QCD have a resonance near threshold ?
- Why care ?
 - Concerns the nonperturbative domain of QCD
 - Quark and gluon degrees of freedom useless there
 - ➔ Understanding very poor, pattern of energy levels ?
 - Lowest resonance: σ ? ρ ?
- Resonance \leftrightarrow pole on second sheet
 - Poles are universal
 - Pole position is unambiguous, even if width is large
 - Where is the pole closest to the origin ?

$f_0(600)$

or σ

$I^G(J^{PC}) = 0^+(0^{++})$

A REVIEW GOES HERE – Check our WWW List of Reviews

$f_0(600)$ T-MATRIX POLE \sqrt{s}

Note that $\Gamma \approx 2 \operatorname{Im}(\sqrt{s}_{\text{pole}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400–1200)–i(300–500) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
(541 ± 39)– i (252 ± 42)	1 ABLIKIM	04A BES2	$J/\psi \rightarrow \omega \pi^+ \pi^-$
(528 ± 32)– i (207 ± 23)	2 GALLEGO	04 RVUE	Compilation
(440 ± 8)– i (212 ± 15)	3 PELAEZ	04A RVUE	$\pi \pi \rightarrow \pi \pi$
(533 ± 25)– i (247 ± 25)	4 BUGG	03 RVUE	
532 – i 272	BLACK	01 RVUE	$\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$
(470 ± 30)– i (295 ± 20)	5 COLANGELO	01 RVUE	$\pi \pi \rightarrow \pi \pi$
(535 ⁺⁴⁸ ₋₃₆)– i (155 ⁺⁷⁶ ₋₅₃)	6 ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon \pi \pi$
610 ± 14 – i 620 ± 26	7 SUROVTSEV	01 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
(558 ⁺³⁴ ₋₂₇)– i (196 ⁺³² ₋₄₁)	ISHIDA	00B	$p \bar{p} \rightarrow \pi^0 \pi^0 \pi^0$
445 – i 235	HANNAH	99 RVUE	π scalar form factor
(523 ± 12)– i (259 ± 7)	KAMINSKI	99 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, \sigma \sigma$
442 – i 227	OLLER	99 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
469 – i 203	OLLER	99B RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
445 – i 221	OLLER	99C RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, \eta \eta$
(1530 ⁺⁹⁰ ₋₂₅₀)– i (560 ± 40)	ANISOVICH	98B RVUE	Compilation
420 – i 212	LOCHER	98 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
(602 ± 26)– i (196 ± 27)	8 ISHIDA	97	$\pi \pi \rightarrow \pi \pi$
(537 ± 20)– i (250 ± 17)	9 KAMINSKI	97B RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, 4\pi$
470 – i 250	10,11 TORNQVIST	96 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, K \pi,$ $\eta \pi$
~(1100 – i 300)	AMSLER	95B CBAR	$\bar{p} p \rightarrow 3\pi^0$
400 – i 500	11,12 AMSLER	95D CBAR	$\bar{p} p \rightarrow 3\pi^0$
1100 – i 137	11,13 AMSLER	95D CBAR	$\bar{p} p \rightarrow 3\pi^0$
387 – i 305	11,14 JANSSEN	95 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
525 – i 269	15 ACHASOV	94 RVUE	$\pi \pi \rightarrow \pi \pi$
(506 ± 10)– i (247 ± 3)	KAMINSKI	94 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
370 – i 356	16 ZOU	94B RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
408 – i 342	11,16 ZOU	93 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
870 – i 370	11,17 AU	87 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
470 – i 208	18 BEVEREN	86 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, \eta \eta, \dots$
(750 ± 50)– i (450 ± 50)	19 ESTABROOKS	79 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
(660 ± 100)– i (320 ± 70)	PROTOPOP...	73 HBC	$\pi \pi \rightarrow \pi \pi, K \bar{K}$
650 – i 370	20 BASDEVANT	72 RVUE	$\pi \pi \rightarrow \pi \pi$

Model independent determination of the pole

- All of the results quoted by the PDG are obtained by
 - (a) parametrizing the data for real values of s
 - (b) continuing this parametrization analytically in s
- ⇒ Result is sensitive to the parametrization used

Model independent determination of the pole

- All of the results quoted by the PDG are obtained by
 - (a) parametrizing the data for real values of s
 - (b) continuing this parametrization analytically in s
- ⇒ Result is sensitive to the parametrization used
- We found a model independent method:
 1. Poles on second sheet are zeros on first sheet
 2. The Roy equations are valid for complex values of s , in a limited region of the first sheet
- ⇒ Exact representation of the partial waves in terms of observable quantities, valid for complex values of s

Model independent determination of the pole


- All of the results quoted by the PDG are obtained by
 - (a) parametrizing the data for real values of s
 - (b) continuing this parametrization analytically in s
- ⇒ Result is sensitive to the parametrization used
- We found a model independent method:
 1. Poles on second sheet are zeros on first sheet
 2. The Roy equations are valid for complex values of s , in a limited region of the first sheet
- ⇒ Exact representation of the partial waves in terms of observable quantities, valid for complex values of s
- 3. Can evaluate this representation to good precision and determine the zeros numerically

Pole on second sheet \leftrightarrow zero on first sheet

- $S_0^0(s) = \eta_0^0(s) \exp 2i\delta_0^0(s)$

$S_0^0(s)$ is analytic in the cut plane

s-plane



Pole on second sheet \leftrightarrow zero on first sheet

- $S_0^0(s) = \eta_0^0(s) \exp 2i\delta_0^0(s)$


$S_0^0(s)$ is analytic in the cut plane

- For $0 < s < 4M_\pi^2$, $S_0^0(s)$ is real

$\Rightarrow S_0^0(s^\star) = S_0^0(s)^\star$

x in elastic interval: $S_0^0(x \pm i\epsilon) = \exp \pm 2i\delta_0^0(x)$


s-plane



Pole on second sheet \leftrightarrow zero on first sheet

- $S_0^0(s) = \eta_0^0(s) \exp 2i\delta_0^0(s)$

$S_0^0(s)$ is analytic in the cut plane



- For $0 < s < 4M_\pi^2$, $S_0^0(s)$ is real

$$\Rightarrow S_0^0(s^\star) = S_0^0(s)^\star$$

x in elastic interval: $S_0^0(x \pm i\epsilon) = \exp \pm 2i\delta_0^0(x)$

- Second sheet is reached by continuation across the elastic interval of the right hand cut

$$S_0^0(x - i\epsilon)^{II} = S_0^0(x + i\epsilon) = 1/S_0^0(x - i\epsilon)$$

Analyticity \Rightarrow
$$S_0^0(s)^{II} = 1/S_0^0(s)$$
 valid $\forall s$

Pole in $S_0^0(s)^{II}$ \iff zero in $S_0^0(s)$

Roy equation for the isoscalar S -wave

$$S_0^0(s) = 1 + 2 i \rho t_0^0(s) \quad \rho = \sqrt{1 - 4 M_\pi^2/s}$$

Roy equation for the isoscalar S -wave

$$S_0^0(s) = 1 + 2 i \rho t_0^0(s) \quad \rho = \sqrt{1 - 4 M_\pi^2 / s}$$

$$\begin{aligned} t_0^0(s) = & a + (s - 4M_\pi^2) b + \int_{4M_\pi^2}^{\infty} ds' \left\{ K_0(s, s') \operatorname{Im} t_0^0(s') \right. \\ & \left. + K_1(s, s') \operatorname{Im} t_1^1(s') + K_2(s, s') \operatorname{Im} t_0^2(s') \right\} \\ & + \text{higher partial waves} \end{aligned}$$

Roy equation for the isoscalar S -wave

$$S_0^0(s) = 1 + 2i\rho t_0^0(s) \quad \rho = \sqrt{1 - 4M_\pi^2/s}$$

$$\begin{aligned} t_0^0(s) = & a + (s - 4M_\pi^2) b + \int_{4M_\pi^2}^{\infty} ds' \left\{ K_0(s, s') \operatorname{Im} t_0^0(s') \right. \\ & \left. + K_1(s, s') \operatorname{Im} t_1^1(s') + K_2(s, s') \operatorname{Im} t_0^2(s') \right\} \\ & + \text{higher partial waves} \end{aligned}$$

- The subtraction constants are determined by a_0^0, a_0^2 :
 $a = a_0^0, \quad b = (2a_0^0 - 5a_0^2)/(12M_\pi^2)$

Roy equation for the isoscalar S -wave

$$S_0^0(s) = 1 + 2i\rho t_0^0(s) \quad \rho = \sqrt{1 - 4M_\pi^2/s}$$

$$\begin{aligned} t_0^0(s) = & a + (s - 4M_\pi^2) b + \int_{4M_\pi^2}^{\infty} ds' \left\{ K_0(s, s') \operatorname{Im} t_0^0(s') \right. \\ & \left. + K_1(s, s') \operatorname{Im} t_1^1(s') + K_2(s, s') \operatorname{Im} t_0^2(s') \right\} \\ & + \text{higher partial waves} \end{aligned}$$

- The subtraction constants are determined by a_0^0, a_0^2 :
 $a = a_0^0, \quad b = (2a_0^0 - 5a_0^2)/(12M_\pi^2)$
- The kernels are elementary functions, e.g.

$$K_0(s, s') = \underbrace{\frac{1}{\pi(s' - s)}}_{r.h.cut} + \underbrace{\frac{2 \ln\{(s + s' - 4M_\pi^2)/s'\}}{3\pi(s - 4M_\pi^2)} - \frac{5s' + 2s - 16M_\pi^2}{3\pi s'(s' - 4M_\pi^2)}}_{l.h.cut}$$

Roy equation for the isoscalar S -wave

$$S_0^0(s) = 1 + 2i\rho t_0^0(s) \quad \rho = \sqrt{1 - 4M_\pi^2/s}$$

$$\begin{aligned} t_0^0(s) = & a + (s - 4M_\pi^2) b + \int_{4M_\pi^2}^{\infty} ds' \left\{ K_0(s, s') \operatorname{Im} t_0^0(s') \right. \\ & \left. + K_1(s, s') \operatorname{Im} t_1^1(s') + K_2(s, s') \operatorname{Im} t_0^2(s') \right\} \\ & + \text{higher partial waves} \end{aligned}$$

- The subtraction constants are determined by a_0^0, a_0^2 :
 $a = a_0^0, \quad b = (2a_0^0 - 5a_0^2)/(12M_\pi^2)$
- The kernels are elementary functions, e.g.

$$K_0(s, s') = \underbrace{\frac{1}{\pi(s'-s)}}_{r.h.cut} + \underbrace{\frac{2\ln\{(s+s'-4M_\pi^2)/s'\}}{3\pi(s-4M_\pi^2)}}_{l.h.cut} - \underbrace{\frac{5s'+2s-16M_\pi^2}{3\pi s'(s'-4M_\pi^2)}}$$
- Left hand cut is essential for convergence:
 $K_0(s, s') \sim 1/s'^3$ for large s'


Domain of validity of Roy equations

- Roy derived his equations from first principles for real energies in the interval $-4M_\pi^2 < s < 60M_\pi^2$

Domain of validity of Roy equations

- Roy derived his equations from first principles for real energies in the interval $-4M_\pi^2 < s < 60M_\pi^2$
- Equations are valid for complex s in a limited region of the first sheet


Caprini, Colangelo & L. 2005



Domain of validity of Roy equations

- Roy derived his equations from first principles for real energies in the interval $-4M_\pi^2 < s < 60M_\pi^2$
- Equations are valid for complex s in a limited region of the first sheet

Caprini, Colangelo & L. 2005



→ Exact representation for $S_0^0(s)$ in this region
Do not need to parametrize the amplitude

Evaluation of the pole position

- Insert our solutions of the Roy equations

For the central solution, $S_0^0(s)$ has two pairs of zeros in the region of validity of the representation:

$$s = (6.2 \pm i 12.3) M_\pi^2 \quad \sigma$$

$$s = (51.4 \pm i 1.4) M_\pi^2 \quad f_0(980)$$


Evaluation of the pole position

- Insert our solutions of the Roy equations

For the central solution, $S_0^0(s)$ has two pairs of zeros in the region of validity of the representation:

$$s = (6.2 \pm i 12.3) M_\pi^2 \quad \sigma$$

$$s = (51.4 \pm i 1.4) M_\pi^2 \quad f_0(980)$$




Evaluation of the pole position

- Insert our solutions of the Roy equations

For the central solution, $S_0^0(s)$ has two pairs of zeros in the region of validity of the representation:

$$s = (6.2 \pm i 12.3) M_\pi^2 \quad \sigma$$

$$s = (51.4 \pm i 1.4) M_\pi^2 \quad f_0(980)$$



⇒ 1. Lowest resonance of QCD
has vacuum quantum numbers


Evaluation of the pole position

- Insert our solutions of the Roy equations

For the central solution, $S_0^0(s)$ has two pairs of zeros in the region of validity of the representation:

$$s = (6.2 \pm i 12.3) M_\pi^2 \quad \sigma$$

$$s = (51.4 \pm i 1.4) M_\pi^2 \quad f_0(980)$$



- ⇒ 1. Lowest resonance of QCD has vacuum quantum numbers
2. Pole on lower half of sheet II occurs in vicinity of

$$\begin{aligned} m_\sigma &= 441 - i 272 \text{ MeV} \\ &= M_\sigma - \frac{i}{2} \Gamma_\sigma \end{aligned}$$

Error analysis

- Results depend on phenomenological input used when solving the Roy equations, subject to uncertainties
Can follow error propagation explicitly
- Pole position of σ mainly depends on 3 input variables:
 $a_0^0, a_0^2, \delta_A \equiv \delta_0^0(800 \text{ MeV})$
 - Substantial uncertainties in phenomenology of δ_A
 - Use the range: $\delta_A = 82.3^\circ {}^{+10^\circ}_{-4^\circ}$

Error analysis

- Noise from remaining input variables is very small:

$$m_\sigma^0 = (441 \pm 4) - i(272 \pm 6) \text{ MeV}$$

Error analysis

- Noise from remaining input variables is very small:

$$m_\sigma^0 = (441 \pm 4) - i(272 \pm 6) \text{ MeV}$$

- Values of a_0^0, a_0^2, δ_A are crucial:

$$\Delta m_\sigma = (-2.4 + i 3.8) \frac{a_0^0 - 0.22}{0.005}$$

$$+ (0.8 - i 4.0) \frac{a_0^2 + 0.0444}{0.001}$$

$$+ (5.3 + i 3.3) \frac{\delta_A - 82.3}{3.4}$$

numbers in MeV

Error analysis

- Noise from remaining input variables is very small:

$$m_\sigma^0 = (441 \pm 4) - i(272 \pm 6) \text{ MeV}$$


- Values of a_0^0, a_0^2, δ_A are crucial:

$$\begin{aligned}\Delta m_\sigma = & (-2.4 + i 3.8) \frac{a_0^0 - 0.22}{0.005} \\ & + (0.8 - i 4.0) \frac{a_0^2 + 0.0444}{0.001} \\ & + (5.3 + i 3.3) \frac{\delta_A - 82.3}{3.4}\end{aligned}\quad \text{numbers in MeV}$$



- Final result: insert the predictions for a_0^0, a_0^2 , use the phenomenological range for δ_A and add errors up:

$$m_\sigma = 441 {}^{+16}_{-8} - i 272 {}^{+9}_{-13} \text{ MeV}$$

Comparison with compilation of PDG




Vicinity of the pole



Results for $\text{Re}[m_\sigma]$ and $\text{Im}[m_\sigma]$ are strongly correlated

Ignore the theoretical predictions for a_0^0, a_0^2

- Replace the low energy theorems for a_0^0, a_0^2 by the experimental results from E865, DIRAC and NA48
- $a_0^0, a_0^2 \in$ universal band



Why are our errors so incredibly small ?

- The σ occurs at low energies
- At low energies, the subtraction term dominates

$$t_0^0(s) \simeq a_0^0 + (2a_0^0 - 5a_0^2) \frac{(s - 4M_\pi^2)}{12M_\pi^2}$$

Why are our errors so incredibly small ?

- The σ occurs at low energies
- At low energies, the subtraction term dominates

$$t_0^0(s) \simeq a_0^0 + (2a_0^0 - 5a_0^2) \frac{(s - 4M_\pi^2)}{12M_\pi^2}$$


Insert low energy theorem for a_0^0, a_0^2

⇒ Roy equation reduces to Weinberg formula

$$t_0^0(s) \simeq \frac{(2s - M_\pi^2)}{32\pi F_\pi^2}$$

Dispersion integrals only represent a correction

At low energies, the subtraction term dominates



$$s = (0.41 \pm 0.06) M_\pi^2 \text{ Adler zero}$$

$$s = (6.2 - i 12.3) M_\pi^2 \text{ pole from } \sigma$$

Estimate pole position on back of an envelope

- Approximate $t_0^0(s)$ with the Weinberg formula

$$t_0^0(s) = \frac{(2s - M_\pi^2)}{32\pi F_\pi^2}$$

Where are the zeros of $S_0^0(s)$ in this approximation ?

Estimate pole position on back of an envelope

- Approximate $t_0^0(s)$ with the Weinberg formula

$$t_0^0(s) = \frac{(2s - M_\pi^2)}{32\pi F_\pi^2}$$

Where are the zeros of $S_0^0(s)$ in this approximation ?

$$1 + 2i\sqrt{1 - 4M_\pi^2/s} t_0^0(s) = 0$$

Estimate pole position on back of an envelope

- Approximate $t_0^0(s)$ with the Weinberg formula

$$t_0^0(s) = \frac{(2s - M_\pi^2)}{32\pi F_\pi^2}$$

Where are the zeros of $S_0^0(s)$ in this approximation ?

$$1 + 2i\sqrt{1 - 4M_\pi^2/s} t_0^0(s) = 0$$

⇒ Cubic equation for s

- Pair of complex zeros, $m_\sigma = 365 - i 291$ MeV

Estimate pole position on back of an envelope

- Approximate $t_0^0(s)$ with the Weinberg formula

$$t_0^0(s) = \frac{(2s - M_\pi^2)}{32\pi F_\pi^2}$$

Where are the zeros of $S_0^0(s)$ in this approximation ?

$$1 + 2i\sqrt{1 - 4M_\pi^2/s} t_0^0(s) = 0$$

⇒ Cubic equation for s

- Pair of complex zeros, $m_\sigma = 365 - i 291$ MeV
- Correction from higher orders amounts to

$$\Delta m_\sigma = 76 {}^{+16}_{-8} + i 19 {}^{+9}_{-13} \text{ MeV}$$

For the quantity that counts, the accuracy is modest

Estimate pole position on back of an envelope

- Approximate $t_0^0(s)$ with the Weinberg formula

$$t_0^0(s) = \frac{(2s - M_\pi^2)}{32\pi F_\pi^2}$$

Where are the zeros of $S_0^0(s)$ in this approximation ?

$$1 + 2i\sqrt{1 - 4M_\pi^2/s} t_0^0(s) = 0$$

⇒ Cubic equation for s

- Pair of complex zeros, $m_\sigma = 365 - i 291$ MeV
- Correction from higher orders amounts to

$$\Delta m_\sigma = 76 {}^{+16}_{-8} + i 19 {}^{+9}_{-13} \text{ MeV}$$

For the quantity that counts, the accuracy is modest

- Real zero on sheet II, near $s = 0$ (full amplitude has kinematic singularity: vanishes on sheet II at $s = 0$)

Conclusion

- Low energy pion physics: theory ahead of experiment
 - Precision experiments carried out and under way
 - Lattice makes slow, but steady progress
 - So far all tests confirm the theory
 - Application to pion e.m. form factor, hadronic contributions to muon g-2

→ talk by G. Colangelo

Conclusion

- Low energy pion physics: theory ahead of experiment
 - Precision experiments carried out and under way
 - Lattice makes slow, but steady progress
 - So far all tests confirm the theory
 - Application to pion e.m. form factor, hadronic contributions to muon g-2
- talk by G. Colangelo
- Limitations of our approach
 - Calculations cannot be done on back of an envelope
 - Method only covers low energies
 - So far, only $\pi\pi$ scattering and form factors
 $J/\psi \rightarrow \omega\pi\pi, D \rightarrow 3\pi, \pi K, \kappa, \dots ??$

Conclusion

- Model independent method for analytic continuation
 - The lowest resonance of QCD occurs at
$$M_\sigma = 441 {}^{+16}_{-8} \text{ MeV} \quad \Gamma_\sigma = 544 {}^{+18}_{-25} \text{ MeV}$$
and carries vacuum quantum numbers
 - Crossing symmetry plays an essential role:
Fixes contributions from left hand cut
Ensures fast convergence, low energy dominance
 - Pole occurs at low value of s , closer to left hand cut than to singularities from $K\bar{K}$, $f_0(980)$
 - Result for Γ_σ relies on theory for a_0^2
Experiments concerning a_0^2 would be most welcome